

## Weiterführende Literatur (in alphabetischer Reihenfolge):

- Adli, Baethge et al. 2005: Is dose escalation of antidepressants a rational strategy after a medium-dose treatment has failed? A systematic review. *Eur Arch Psychiatry Clin Neurosci* 255(6): 387-400
- Bajbouj et al. *J of Clinical Psychopharm* 2010
- Bakker et al. 2015
- Baldinger P et al. *J ECT*. 2014 Jun; 30(2): 116-21
- Barnhofer et al., 2009. Mindfulness-based cognitive therapy as a treatment for chronic depression. A preliminary study. *Behav Res Ther*; 47:366-373
- Bartova L., Dold M., Weidenauer A., Kasper S. (2017): Behandlungsresistente Depression, *Clinicum neuropsych* 3/17: 28-34
- Bauer, Severus et al. 2017. Pharmacological treatment of unipolar depressive disorders: summary of WFSBP guidelines. *Int J Psychiatry Clin Pract*: 1-11
- Bewernick et al. *Biol Psychiatry* 2010
- Brakemeier et al., 2015. Overcoming treatment resistance in chronic depression: a pilot study on outcome and feasibility of the cognitive behavioral analysis system of psychotherapy as an inpatient treatment program. *Psychother Psychosom* 84:51–56
- Brent et al., 2008. Switching to another SSRI or to venlafaxine with or without cognitive behavioral therapy for adolescents with SSRI-resistant depression: the TORDIA randomized controlled trial. *JAMA*. 299:901-13.
- Bschor et al *DD: Ärzteblatt*, Heft 45, 7.Nov.2014
- Bschor et al., 2016, *J Clin Psychiatry*
- Cascorbi, 2012. Arzneimittelinteraktionen. *Deutsches Ärzteblatt*, 109:33-34, 546-555.
- Cooper C et al. *Am J Psychiatry*, March 31 2011
- Coplan et al. 2015. Treating comorbid anxiety and depression: Psychosocial and pharmacological approaches. *World J Psychiatry* 5(4): 366-378
- Crossley and Bauer, 2007. Acceleration and augmentation of antidepressants with lithium for depressive disorders: two meta-analyses of randomized, placebo-controlled trials. *J Clin Psychiatry* 68(6): 935-940
- Dekker et al., 2013. What is the best sequential treatment strategy in the treatment of depression? Adding pharmacotherapy to psychotherapy and vice versa? *Psychother Psychosom* 82:89-98
- DGPPN, BÄK, KBV, AWMF, AkdÄ, BpTK, BApK, DAGSHG, DEGAM, DGPM, DGPs and DGRW (2015). S3-Leitlinie/Nationale VersorgungsLeitlinie Unipolare Depression – Langfassung
- Dold, Kautzky et al. 2016, Pharmacological treatment strategies in unipolar depression in European tertiary psychiatric treatment centers - A pharmacoepidemiological cross-sectional multicenter study. *Eur Neuropsychopharmacol* 26(12): 1960-1971
- Dold, Bartova et al. 2017. Dose escalation of antidepressants in unipolar depression: a meta-analysis of double-blind, randomized controlled trials. *Psychotherapy and Psychosomatics*
- Dold M, Kasper S. Evidence-based pharmacotherapy of treatment-resistant unipolar depression. *Int J Psychiatry Clin Pract*. 2017: 1-11
- Eisendrath et al., 2016. A randomized controlled trial of mindfulness-based cognitive therapy for treatment-resistant depression. *Psychother Psychosom* 85(2): 99-110
- Emslie et al., 2010. Treatment of Resistant Depression in Adolescents (TORDIA): week 24 outcomes. *Am J Psych*. 167(7):782-91
- Fonagy et al., 2015. Pragmatic randomized controlled trial of long-term psychoanalytic psychotherapy for treatment-resistant depression: the Tavistock Adult Depression Study (TADS). *World Psychiatry* 14:312–321
- Goodyear et al., 2007. Selective serotonin reuptake inhibitors (SSRIs) and routine specialist care with and without cognitive behaviour therapy in adolescents with major depression: randomized controlled trial. *Br Med J*. ;335:142

- Gören und Tewksbury, 2013. Drug interactions and Polypharmacy. In Polypharmacy in Psychiatry Practice, Volume I: Multiple Medication Use Strategies; Ritsner MS. (Editor), Springer Verlag, Dordrecht, 45-74
- Greiner, 2009
- Hamilton, 1960. A rating scale for depression. Journal of neurology, neurosurgery, and psychiatry 23: 56-62
- Hautzinger et al., 2013. Kognitive Verhaltenstherapie bei Depressionen: Behandlungsanleitungen und Materialien. Beltz, Weinheim
- Hedrick et al., Cochrane 2007;(3);CD004851
- Hiemke et al., 2017. Consensus Guidelines for Therapeutic Drug Monitoring in Neuropsychopharmacology: Update 2017. Pharmacopsychiatry, epub ahead of print 14 Sep 2017, DOI: 10.1055/s-0043-116492
- Höflich et al. Neuroimage 2013; 79:30-41
- Hollon et al., 2014. Effect of cognitive therapy with antidepressant medications vs antidepressants alone on the rate of recovery in major depressive disorder: a randomized clinical trial. JAMA Psychiatry 71:1157–1164
- Holsboer-Trachsler et al., 2016. Die Akutbehandlung depressiver Episoden. Swiss medical forum - schweizerisches Medizin-Forum 16: 716-724
- Hovstadius und Petersson, 2013. The impact of increasing polypharmacy on prescribed drug expenditure – a register-based study in Sweden 2005-2009. Health Policy 109, 166-174
- Johnell und Klarin, 2007. The relationship between number of drugs and potential drug-drug interactions in the elderly: a study of over 600.000 elderly patients from the Swedish Prescribed Drug Register. Drug Saf 30, 911-918 (2007)
- Kasper und Akimova, 2013. The Role of Atypical Antipsychotics in Inadequate-response and Treatment-resistant Depression. Treatment-resistant depression. S. Kasper and S. Montgomery. Chichester: Wiley-Blackwell: pp: 107-128
- Kasper, Bach et al. 2011
- Kasper et al., 2012. Depression – Medikamentöse Therapie. Clinicum neuropsy. Sonderausgabe November 2012
- Keller et al., 2000. A comparison of nefazodone, the cognitive behavioral-analysis system of psychotherapy, and their combination for the treatment of chronic depression. N Engl J Med 342:1462–1470
- Kennedy et al. Am J Psychiatry 2011
- Klesse C, Berger M et al., 2010. Evidenzbasierte Psychotherapie der Depression. Die S3-Leitlinie unipolare Depression. Psychotherapeut. 247–263
- Köhler et al., 2016. Überwindung der Therapieresistenz bei chronischer Depression. Die Bedeutung der stationären Psychotherapie. Nervenarzt 87:701–707
- Kok RM et al. Acta Psychiatr Scand. 2009 April; 119(4):274–81
- Kolbinger HM et al. Human Psychopharmacology 10:305-310
- Konstantinidis A et al. Int J Neuropsychopharm; epub ahead of print 17 May 2011
- Kraus C et al., 2017. Administration of ketamine for unipolar and bipolar depression. International Journal of Psychiatry in Clinical Practice 21:2-12
- Lanzenberger R et al. Molecular Psychiatry 2013; 18(1): 93-100
- Leucht, Hierl et al., 2012. Putting the efficacy of psychiatric and general medicine medication into perspective: review of meta-analyses. Br J Psychiatry 200(2): 97-106
- Levkovitz et al. 2015;
- Lingham and Scott, Acta Psych Scand. 2002; 105:164–173
- Malone et al. Biol Psychiatry 2009; 65:267-75
- March, Silver et al., 2009. The Treatment for Adolescents With Depression Study (TADS): outcomes over 1 year of naturalistic follow-up. Am J Psychiatry 2009;166:1141–9

- McKnight, Adida et al., 2012. Lithium toxicity profile: a systematic review and meta-analysis. *Lancet* 379(9817): 721-728
- Montgomery und Asberg, 1979. A new depression scale designed to be sensitive to change. *Br J Psychiatry*: 134:382-389
- Nakagawa et al., 2017. Effectiveness of supplementary cognitive-behavioral therapy for pharmacotherapy-resistant depression: A randomized controlled trial. *J Clin Psychiatry* doi: 10.4088/JCP.15m10511
- Nelson und Papakostas, 2009. Atypical antipsychotic augmentation in major depressive disorder: a meta-analysis of placebo-controlled randomized trials. *Am J Psychiatry* 166(9): 980-991
- Neuner-Jehle, 2013. Less is more – how to prevent polypharmacy?. *Praxis (Bern 1994)* 102, 21-27
- O’Readon et al. *Biol Psychiatry* 2007, George et al. *Arch Gen Psychiatry* 2010, Avers et al. *Biol Psychiatry* 2006
- Parker und Graham, 2015. Determinants of treatment-resistant depression: The salience of benzodiazepines. *J Nerv Ment Dis* 203: 659-663
- Rush et al. *Biol Psychiatry* 2000 und 2005
- Sackeim et al. *Arch Gen Psychiatry*. 2000;57:425–434
- Sakolsky et al., 2011. Antidepressant exposure as a predictor of clinical outcomes in the Treatment of Resistant Depression in Adolescents (TORDIA) study. *J Clin Psych*. 31(1):92-7
- Sartorius et al. *Biol Psychiatry* 2010; 15:67
- Sandson, 2003. *Drug Interactions Casebook. The Cytochrome P450 System and Beyond.* American Psychiatric Publishing, Inc. 2003, Arlington, United States of America.
- Schlaepfer et al. *WFSBP Guidelines on Brain Stimulation Treatments in Psychiatry. The World Journal of Biological Psychiatry* 2010; 11:2-18
- Schosser, Serretti et al., 2012. European Group for the Study of Resistant Depression (GSRD)- where have we gone so far: review of clinical and genetic findings." *Eur Neuropsychopharmacol* 22(7): 453-468
- Schramm E et al., 2017. Effect of disorder-specific vs nonspecific psychotherapy for chronic depression: A randomized clinical trial. *JAMA Psychiatry* 74(3):233-242
- Silberman und Weiss, 2016. Treatment-resistant depression or misdiagnosed anxiety disorder? *J Nerv Ment Dis* 204: 67
- Souery et al., 1999. Treatment resistant depression: methodological overview and operational criteria. *Eur Neuropsychopharmacol* 9:83-91.
- Souery D et al., 2007. Clinical factors associated with treatment resistance in major depressive disorder: results from a European multicenter study. *J Clin Psychiatry*. 68:7,1062–70
- Souery D et al. Switching antidepressant class does not improve response or remission in treatment-resistant depression. *J Clin Psychopharmacol* 2011; 31(4):512–516
- Spindelegger, Papageorgiou, et al., 2014. Cardiovascular adverse reactions during antidepressant treatment: a drug surveillance report of German-speaking countries between 1993 and 2010. *Int J Neuropsychopharmacol* 18(4).
- Thase ME, Rush AJ., 1997. When at first you don't succeed: sequential strategies for antidepressant nonresponders. *J Clin Psychiatry*. 58 Suppl 13:23–29
- Trivedi et al., 2011. Examination of the utility of psychotherapy for patients with treatment resistant depression: a systematic review. *J Gen Intern Med*. 26(6): 643-650
- Van Straten et al., 2014. Stepped care treatment delivery for depression: a systematic review and meta-analysis. *Psychol Med* 45(2):1–16
- Wiersma et al., 2014. The effectiveness of the cognitive behavioral analysis system of psychotherapy for chronic depression: a randomized controlled trial. *Psychother Psychosom* 83:263–269
- Yoshino et al. *Psychiatry Clin Neurosci*. 2009; 63(5):652–657