

Literatur zur Expertise „Psychosen im Alter“

- Albert SG, Grossberg GT, Thaipisuttikul PJ, Scouby J, Green E. A typical antipsychotics and the risk of diabetes in an elderly population in long-term care: a retrospective nursing home chart review study. *J Am Med Dir Assoc* 2009 Feb; 10(2) :115–9
- Ciranni MA, Kearney TE, Olson KR. Comparing acute toxicity of first- and second-generation antipsychotic drugs: A 10-year, retrospective cohort study. *J Clin Psychiatry* 2009 Jan 13
- Edwards K et al. Efficacy and Safety of Galantamine in Patients with Dementia with Lewy Bodies: A 24-Week Open-Label Study. *Dement Geriatr Cogn Disord* 2007; 23: 401–405
- Howard R et al. International Late-onset Schizophrenia Group. Late-onset schizophrenia and very-late-onset schizophrenia like psychosis: an international consensus. *Am J Psych* 2000; 157: 172–178
- Howland RH. Risks and benefits of antipsychotic drugs in elderly patients with dementia. *J Psychosoc Nurs Ment Health Serv* 2008 Nov; 46(11): 19–23
- Jeste DV, Finkel SI. Psychosis of Alzheimer's disease and related dementias: diagnostic criteria for a distinct syndrome. *Am J Geriatr Psychiatry* 2000
- Jeste DV, Twamley EW. Understanding and managing psychosis in late life. *Psych Times* 2003; 20(3): 19–22
- Karim S, Byrne EJ. Treatment of psychosis elderly people. *Br J Psych* 2005; 11: 286–296
- Kassabian S. Antipsychotic therapy for dementia in elderly patients. *Arch Intern Med* 2009 Jan 12; 169(1): 87–8
- Keck PE et al. Placebo-controlled, double-blind study of the efficacy and safety of aripiprazole in patients with acute bipolar mania. *Am J Psych* 2003; 160: 1651–1658
- Leucht S, Coves C. Second generation versus first generation antipsychotic drugs for schizophrenia: meta analysis. *Lancet* 2009; 373: 31–41
- Mintzer J, Targum SD. Psychosis in elderly patients: classification and pharmacotherapy. *Psych Neurol* 2003; 16(4): 199–206
- Pirmohamed et al. Adverse drug reactions as cause of admission to hospital: prospective analysis of 18820 patients. *Brit.Med.J* 2004; 329: 15–19
- Schneeweiss S, Setoguchi S, Brookhart MA, Kaci L, Wang PS. Assessing Residual Confounding of the Association between Antipsychotic Medications and Risk of Death using Survey Data, *CNS Drugs* 2009; 23(2): 171–80
- Spar JE, LaRue A. Concise Guide to Geriatric Psychiatry. 3rd ed. Washington, Psychiatric Publishing 2006
- Tadger S. et al. Symptomatic remission in elderly schizophrenia patients treated with long-acting risperidone. *Int Psychogeriatrics* 2008; 1–6
- Tohen M et al. Olanzapine versus divalproex sodium for the treatment of acute mania and maintenance of remission: a 47-week study. *Am J Psych* 2003;169: 1263–1271
- Uchida H, Mamo DC, Mulsant BH, Pollock BG, Kapur S. Increased antipsychotic sensitivity in elderly patients: Evidence and mechanisms. *J Clin Psychiatry* 2008 Dec 16; pii: ej08r04171
- Webster J, Grossberg GT. Late life psychotic symptoms. *Am J Geriatr Psych* 1998; 6(3): 196–202
- Wylie ME, Mulsant BH, Pollock BG et al. Age at onset in geriatric bipolar disorder and clinical presentation and treatment outcomes in an inpatient sample. *Am J Geriatr Psych* 1999; 7(1): 77–83
- Yatham LN. Quetiapine versus Placebo in combination with lithium or divalproex for the treatment of bipolar mania. *J Clin Psychopharmacol.* 2004; 24(6): 599–606
- Yatham LN et al. Mood stabilisers plus risperidone or placebo in the treatment of acute mania: an international, double-blind, randomised, controlled trial. *Br J Psychiatry* 2003; 182: 141–147
- Yatham LN et al. Canadian Network for Mood and Anxiety Treatments (CANMAT) guidelines for the management of patients with bipolar disorder: consensus and controversies. *Bipolar Disorders* 2005; 7 (Suppl. 3): 5–69